

## **Product Datasheet**

Flavivirus group antibody (orb357996)



**Description**nts. Recombinant monoclonal antibody to Flavivirus

## www.biorbyt.com

Mock (Media) 2\*Ab only control ZIKV (MOI 0.2

E protein Merge

Immunofluorescense analysis of Vero cell...

Species/Host

Mouse

Reactivity

Virus

Conjugation

Unconjugated

Tested

ELISA, FC, IF, IHC-P, WB

**Applications** 

- , -, , - ,

Immunogen

Dengue Virus type 2 antigens.

**Target** 

Flavivirus group antigen

**Preservatives** 

PBS with 0.02% Proclin 300.

Concentration

1 mg/ml

Storage

Store at 4°C for up to 3 months. For longer

storage, aliquot and store at -20°C.

Note

For research use only

**Application notes** 

This antibody binds to flavivirus group antigen, protein E. It can be used as an anti-Dengue virus antibody, anti-West Nile virus antibody, anti-Japanese Encephalitis, anti-Yellow Fever Virus or anti-Zika Virus antibody (Aubry et al. 2016) to identify cells infected with these flaviviridae. It binds to the fusion loop at the extremity of domain II of E protein from all four serotypes and prevents syncytia formation (Summers, 1989). The epitope is highly conserved amongst

flaviviridae and has been functionally analyzed in detail by Crill and Chang 2004 (PMID: 15564505). Previous studies have used acetone- (Henchal et al. 1982) or methanol-fixed slides (Moreland & Tay, 2010). Please note that binding of this antibody has been reported to be sensitive to reduction and Western Blots should be performed under non-reducing conditions (Lai et al. 2008). A version of 4G2 directly conjugaed to APC was used by Quicke and co-workers (Quicke, 2016) to detect Zika virus in human placental

macrophages by flow-cytometry (Quicke 2016).

Ramaiah et al. (2016) used Ab00230-2.0 at a 1:200 dilution to stain for Zika virus in methanol-

fixed Vero cells 48 h post-infection.

Isotype

IgG

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Clonality

Monoclonal

Purity

Purified