



## **Product Datasheet**

**TWEAK antibody (orb669765)** 



## www.biorbyt.com

**Description**nts. Mouse monoclonal antibody to TWEAK

Species/Host Rat

Reactivity Mouse

**Conjugation** Unconjugated

Tested

**Applications** 

Blocking, FC

Immunogen This antibody was raised by immunising SD rats

with CD8-mTWEAK, a fusion of the human CD8alpha extracellular region to murine

TWEAK.

Target TWEAK

**Preservatives** PBS with 0.02% Proclin 300.

**Concentration** 1 mg/ml

**Storage** Store at 4°C for up to 3 months. For longer

storage, aliquot and store at -20°C.

**Note** For research use only

**Application notes** The specificity of this antibody has been

reacts specifically with mTWEAK-transfected L5178Y cells, and not with parental L5178Y cells or cells transfected with related TNF family members (Nakayama et al, 2003). Additionally, this antibody has been used to assess TWEAK cell surface expression in flow cytometric analysis (Nakayama et al, 2003). This antibody has been shown to block CD8-mTWEAK cytotoxicity against L5178Y cells over-expressing murine Fn14, the TWEAK receptor (Nakayama et al, 2003). When administered to collagen-induced arthritis (CIA) mice, this

confirmed in flow cytometric analysis, where it

(Nakayama et al, 2003). When administered to collagen-induced arthritis (CIA) mice, this blocking antibody has been shown to have anti-inflammatory and antiangiogenic effects, with associated improvements to both the clinical and histological manifestations of CIA (Kamata et al, 2006). Additionally, administration of this antibody to denervated mice attenuates denervation-induced skeletal muscle loss (Mittal et al, 2010). Finally, administration of this antibody to CD163-/- mice subject to hind limb ischaemia inhibits the increased perfusion resulting from CD163 deficiency (Akahori et al,

2015).

Isotype IgG1

Biorbyt Ltd.

Biorbyt LLC.