

## **Product Datasheet**

Rat IgG2a Isotype Control antibody (APC) (orb699899)



**Description**nts. Rat monoclonalantibody against Rat IgG2a Isotype

## www.biorbyt.com

**Conjugation** APC

**Tested** FC **Applications** 

**Preservatives** Stabilizing phosphate buffered saline (PBS), pH 7.4,

15 mM sodium azide

**Concentration** 0.1 mg/ml

**Storage** Store at 2-8°C. Protect from prolonged exposure to

light. Do not freeze.

**Note** For research use only

**Application notes** Negative control: The reagent is intended as an

isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg/ml of incubation mixture), change the

conditions of your experiment to reduce the

background.

**Isotype** Rat IgG2a


**Clonality** Monoclonal

**Purity** Purified antibody is conjugated with activated

allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

**Dilution Range** Negative control: The reagent is intended as an

isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background



Flow cytometry surface nonspecific stain...