



Product Datasheet

Human GYPB protein (orb705114)

Description	Recombinant Human Glycophorin-B(GYPB)
Endotoxins	Not test.
Preservatives	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.
Form/Appearance	Liquid or Lyophilized powder
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Тад	N-terminal 10xHis-tagged
Note	For research use only
Application notes	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Application notes Protein Sequence	bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of
	bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. LSTTEVAMHTSTSSSVTKSYISSQTNGETGQLVHRFTVPAPVVIILIILCVMAGIIGT
Protein Sequence	bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. LSTTEVAMHTSTSSSVTKSYISSQTNGETGQLVHRFTVPAPVVIILIILCVMAGIIGT ILLISYSIRRLIKA
Protein Sequence Purity	bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. LSTTEVAMHTSTSSSVTKSYISSQTNGETGQLVHRFTVPAPVVIILIILCVMAGIIGT ILLISYSIRRLIKA Greater than 85% as determined by SDS-PAGE.
Protein Sequence Purity Source	bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. LSTTEVAMHTSTSSSVTKSYISSQTNGETGQLVHRFTVPAPVVIILIILCVMAGIIGT ILLISYSIRRLIKA Greater than 85% as determined by SDS-PAGE. in vitro E.coli expression system

Biorbyt Ltd

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: **info@biorbyt.com** | Phone: **+44 (0) 1223 859-353** | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558