

Product Datasheet

H4 K8Ac; K12Ac; K16Ac antibody (orb758977)

Description

Mouse monoclonal antibody to H4 K8Ac;

Species/Host	Mouse
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, IF, IHC, WB
Immunogen	KM-2 antibody was derived from an MRL/lpr lupus mouse.
Target	H4 K8Ac; K12Ac; K16Ac
Preservatives	PBS with 0.02% Proclin 300.
Concentration	1 mg/ml
Storage	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.
Note	For research use only
Application notes	<p>The antibody KM-2 shows significant increase in activity with histone H4 when it is acetylated on lysines 8, 12 and 16 which are hyperacetylated in conditions, such as lupus (SLE); and thus KM-2 can be used in the detection of the apoptosis-induced histone modifications characteristic of lupus and in the analysis of its pathogenesis (Dieker et al., 2007; pmid: 17530637). This antibody was successfully used to identify both Histone H4 as well as H2A in the extracts of Jurkat cells via Western blot, showing considerable increase in activity in cells with induced apoptosis leading to hyperacetylation (Dieker et al., 2007; pmid: 17530637). Similar results were achieved for the immunofluorescence staining of Jurkat cells. The enhanced reactivity of KM-2 with apoptotic H4 and apoptotic H2A was not Jurkat cell specific and was also observed for the monocytic U937 cell line which indicated its general applicability (Dieker et al., 2007; pmid: 17530637). Analogous results were also achieved during the ELISA analysis of plasma samples of (diseased) MRL/lpr lupus mice, with the visible activity enhancement in the sick, apoptotic</p>

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709, United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558