



Cleaved-PARP-1 (D214) rabbit pAb

Cat#: orb763946 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Cleaved-PARP-1 (D214) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

Recommended dilutions WB 1:500-2000, IF 1:50-300, IHC 1:50-300

Immunogen The antiserum was produced against synthesized peptide derived from

human PARP. AA range:165-214

Specificity Cleaved-PARP-1 (D214) Polyclonal Antibody detects endogenous levels of

fragment of activated PARP-1 protein resulting from cleavage adjacent to

D214.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Poly [ADP-ribose] polymerase 1

Gene Name PARP1

Cellular localization Nucleus . Nucleus . Nucleus . Chromosome . Localizes to sites of DNA

damage...

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 24kD

Human Gene ID 142

P09874 **Human Swiss-Prot Number**

PARP1; ADPRT; PPOL; Poly [ADP-ribose] polymerase 1; PARP-1; ADP-ribosyltransferase diphtheria toxin-like 1; ARTD1; NAD(+) ADP-ribosyltransferase 1; ADPRT 1; Poly[ADP-ribose] synthase 1 **Alternative Names**

Background This gene encodes a chromatin-associated enzyme, poly(ADP-

ribosyl)transferase, which modifies various nuclear proteins by poly(ADP-ribosyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes. [provided by

RefSeq, Jul 2008],