



Rb (phospho Ser780) rabbit pAb

Cat#: orb764275 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Rb (phospho Ser780) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human Retinoblastoma around the phosphorylation site of Ser780. AA

range:751-800

Specificity Phospho-Rb (S780) Polyclonal Antibody detects endogenous levels of Rb

protein only when phosphorylated at S780.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Retinoblastoma-associated protein

Gene Name RB1

Cellular localization Nucleus . During keratinocyte differentiation, acetylation by KAT2B/PCAF

is required for nuclear localization. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

Concentration 1 mg/ml

Observed band 106kD


Human Gene ID 5925

Human Swiss-Prot Number P06400


Alternative Names RB1; Retinoblastoma-associated protein; p105-Rb; pRb; Rb; pp110

Background

The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma. [provided by RefSeq, Jul 2008],



Immunofluorescence analysis of Hela cell. 1,Rb (phospho Ser780) Polyclonal Antibody(red) was diluted at 1:200(4° overnight). GFAP Monoclonal Antibody(5C8)(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000(room temperature, 50min).




- + - phospho-peptide
- + non-phospho-peptide

Western Blot analysis of various cells using Phospho-Rb (S780) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).






Explore. Bioreagents.



Western Blot analysis of KB cells using Phospho-Rb (S780) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Retinoblastoma (Phospho-Ser780) Antibody. The picture on the right is blocked with the phospho peptide.