



EGFR (phospho Thr678) rabbit pAb

Cat#: orb764322 (Manual)

For research use only. Not intended for diagnostic use.

| Product Name | EGFR (phospho Thr678) rabbit pAb |
|--------------------------|---|
| Host species | Rabbit |
| Applications | WB;IF;ELISA |
| Species Cross-Reactivity | Human;Mouse;Rat;Monkey |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. |
| Immunogen | The antiserum was produced against synthesized peptide derived from human EGFR around the phosphorylation site of Thr678. AA range:651-700 |
| Specificity | Phospho-EGFR (T678) Polyclonal Antibody detects endogenous levels of EGFR protein only when phosphorylated at T678. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Epidermal growth factor receptor |
| Gene Name | EGFR |
| Cellular localization | Cell membrane ; Single-pass type I membrane protein . Endoplasmic reticulum membrane ; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome membrane. Nucleus . In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:20674546, PubMed:17909029). Endocytosed upon activation by ligand (PubMed:2790960, PubMed:17182860, PubMed:27153536, PubMed:17909029). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055); [Isoform 2]: Secreted. |



www.biorbyt.com

| Purification | The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen. |
|-------------------------|---|
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 175kD |
| Human Gene ID | 1956 |
| Human Swiss-Prot Number | P00533 |
| Alternative Names | EGFR; ERBB; ERBB1; HER1; Epidermal growth factor receptor; Proto- oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1 |
| Background | The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. [provided by RefSeq, Jun 2016], |