

c-Src rabbit pAb**Cat#: orb764939 (Manual)**


For research use only. Not intended for diagnostic use.

| | |
|---------------------------------|---|
| Product Name | c-Src rabbit pAb |
| Host species | Rabbit |
| Applications | WB;IP;IHC;IF;ELISA |
| Species Cross-Reactivity | Human;Mouse;Rat |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. IP 1:50-200 ELISA: 1/5000. Not yet tested in other applications. |
| Immunogen | The antiserum was produced against synthesized peptide derived from human Src. AA range:487-536 |
| Specificity | c-Src Polyclonal Antibody detects endogenous levels of c-Src protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Proto-oncogene tyrosine-protein kinase Src |
| Gene Name | SRC |
| Cellular localization | Cell membrane ; Lipid-anchor . Mitochondrion inner membrane . Nucleus . Cytoplasm, cytoskeleton . Cytoplasm, perinuclear region . Cell junction, focal adhesion . Localizes to focal adhesion sites following integrin engagement (PubMed:22801373). Localization to focal adhesion sites requires myristoylation and the SH3 domain (PubMed:7525268). Colocalizes with PDLIM4 at the perinuclear region, but not at focal adhesions (PubMed:19307596). . |


| | |
|--------------------------------|---|
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 59kD |
| Human Gene ID | 6714 |
| Human Swiss-Prot Number | P12931 |
| Alternative Names | SRC; SRC1; Proto-oncogene tyrosine-protein kinase Src; Proto-oncogene c-Src; pp60c-src; p60-Src |

Background


This gene is highly similar to the v-src gene of Rous sarcoma virus. This proto-oncogene may play a role in the regulation of embryonic development and cell growth. The protein encoded by this gene is a tyrosine-protein kinase whose activity can be inhibited by phosphorylation by c-SRC kinase. Mutations in this gene could be involved in the malignant progression of colon cancer. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using c-Src Polyclonal Antibody diluted at 1:2000



Western Blot analysis of HeLa cells using c-Src Polyclonal Antibody diluted at 1:2000



Western Blot analysis of hela using c-Src Polyclonal Antibody. Antibody was diluted at 1:2000



Western Blot analysis of HELA HEPG2 using c-Src Polyclonal Antibody diluted at 1:2000