



Cyclin E1 rabbit pAb

Cat#: orb764957 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Cyclin E1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat; Pig

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human Cyclin E1. ÂA range:91-140

Specificity Cyclin E1 Polyclonal Antibody detects endogenous levels of Cyclin E1

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name G1/S-specific cyclin-E1

Gene Name CCNE1

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 49kD

Human Gene ID 898

Human Swiss-Prot Number P24864

Alternative Names CCNE1; CCNE; G1/S-specific cyclin-E1

Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in