



Eg5 rabbit pAb

Cat#: orb765106 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Eg5 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human KIF11/Eg5. AA range:892-941

Specificity Eg5 Polyclonal Antibody detects endogenous levels of Eg5 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Kinesin-like protein KIF11

Gene Name KIF11

Cellular localization Cytoplasm . Cytoplasm, cytoskeleton, spindle pole .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 119kD

Human Gene ID 3832

Human Swiss-Prot Number P52732

Alternative Names


KIF11; EG5; KNSL1; TRIP5; Kinesin-like protein KIF11; Kinesin-like protein 1; Kinesin-like spindle protein HKSP; Kinesin-related motor protein Eg5; Thyroid receptor-interacting protein 5; TR-interacting protein 5; TRIP-

Background


kinesin family member 11(KIF11) Homo sapiens This gene encodes a motor protein that belongs to the kinesin-like protein family. Members of this

protein family are known to be involved in various kinds of spindle dynamics. The function of this gene product includes chromosome

positioning, centrosome separation and establishing a bipolar spindle during cell mitosis. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using Eg5 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).




Western Blot analysis of COLO205 cells using Eg5 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).







Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



 $Immunofluorescence\ analysis\ of\ HeLa\ cells,\ using\ KIF11/Eg5\ Antibody.\ The\ picture\ on\ the\ right\ is\ blocked\ with\ the\ synthesized\ peptide.$