

GRP1 rabbit pAb**Cat#: orb765357 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	GRP1 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human GRP1. AA range:351-400
Specificity	GRP1 Polyclonal Antibody detects endogenous levels of GRP1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cytohesin-3
Gene Name	CYTH3
Cellular localization	Cytoplasm, cytosol. Cell membrane ; Peripheral membrane protein . Cell junction, adherens junction . Cell junction, tight junction . Translocates from the cytosol to membranes enriched in phosphatidylinositol 3,4,5-trisphosphate. .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	46kD
Human Gene ID	9265
Human Swiss-Prot Number	O43739
Alternative Names	CYTH3; ARNO3; GRP1; PSCD3; Cytohesin-3; ARF nucleotide-binding site opener 3; Protein ARNO3; General receptor of phosphoinositides 1; Grp1; PH; SEC7 and coiled-coil domain-containing protein 3

Background

This gene encodes a member of the PSCD (pleckstrin homology, Sec7 and coiled-coil domains) family. PSCD family members have identical structural organization that consists of an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP) activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family appear to mediate the regulation of protein sorting and membrane trafficking. This encoded protein is involved in the control of Golgi structure and function, and it may have a physiological role in regulating ADP-ribosylation factor protein 6 (ARF) functions, in addition to acting on ARF1. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using GRP1 Polyclonal Antibody



Western blot analysis of lysate from COS7 cells, using GRP1 antibody.