



GSC2 rabbit pAb

Cat#: orb765359 (Manual)

For research use only. Not intended for diagnostic use.

Product Name GSC2 rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human GSC2. AA range:131-180

Specificity GSC2 Polyclonal Antibody detects endogenous levels of GSC2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Homeobox protein goosecoid-2

Gene Name GSC2

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 25kD

Human Gene ID 2928


Human Swiss-Prot Number 015499

Alternative Names GSC2; GSCL; Homeobox protein goosecoid-2; GSC-2; Homeobox protein

goosecoid-like; GSC-L

Background

Goosecoidlike (GSCL), a homeodomain-containing gene, resides in the critical region for VCFS/DGS on 22q11. Velocardiofacial syndrome (VCFS) is a developmental disorder characterized by conotruncal heart defects, craniofacial anomalies, and learning disabilities. VCFS is phenotypically related to DiGeorge syndrome (DGS) and both syndromes are associated with hemizygous 22q11 deletions. Because many of the tissues and structures affected in VCFS/DGS derive from the pharyngeal arches of the developing embryo, it is believed that haploinsufficiency of a gene involved in embryonic development may be responsible for its etiology. The gene is expressed in a limited number of adult tissues, as well as in early human development. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using GSC2 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).





Explore. Bioreagents.



Western blot analysis of lysates from 293, HepG2, and COLO205 cells, using GSC2 Antibody. The lane on the right is blocked with the synthesized peptide.