



LZK rabbit pAb

Cat#: orb765611 (Manual)

For research use only. Not intended for diagnostic use.

Product Name LZK rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human M3K13. AA range:151-200

Specificity LZK Polyclonal Antibody detects endogenous levels of LZK protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Mitogen-activated protein kinase kinase kinase 13

Gene Name MAP3K13

Cellular localization Cytoplasm . Membrane ; Peripheral membrane protein .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 108kD

Human Gene ID 9175

Human Swiss-Prot Number O43283


Alternative Names MAP3K13; LZK; Mitogen-activated protein kinase kinase kinase 13;

Leucine zipper-bearing kinase; Mixed lineage kinase; MLK

Background The protein encoded by this gene is a member of serine/threonine protein

kinase family. This kinase contains a dual leucine-zipper motif, and has been shown to form dimers/oligomers through its leucine-zipper motif. This kinase can phosphorylate and activate MAPK8/JNK, MAP2K7/MKK7, which suggests a role in the JNK signaling pathway. [provided by RefSeq,


Jul 2008],








Explore. Bioreagents.



Western blot analysis of lysates from HeLa, HUVEC, and HepG2 cells, using M3K13 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using M3K13 antibody.