



Olfactory receptor 1A1 rabbit pAb

Cat#: orb765895 (Manual)


For research use only. Not intended for diagnostic use.

Product Name	Olfactory receptor 1A1 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human OR1A1. AA range:260-309
Specificity	Olfactory receptor 1A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 1A1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 1A1
Gene Name	OR1A1
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
	entoniatography using epitope-specific minutogen.



www.biorbyt.com


Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	8383
Human Swiss-Prot Number	Q9P1Q5
Alternative Names	OR1A1; Olfactory receptor 1A1; Olfactory receptor 17-7; OR17-7; Olfactory receptor OR17-11
Background	olfactory receptor family 1 subfamily A member 1(OR1A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein- mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using Olfactory receptor 1A1 Polyclonal Antibody



www.biorbyt.com



Western blot analysis of lysates from COLO cells, using OR1A1 Antibody. The lane on the right is blocked with the synthesized peptide.