



Olfactory receptor 4K14 rabbit pAb

Cat#: orb765919 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Olfactory receptor 4K14 rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Monkey

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human OR4K14. AA range:261-310

Specificity Olfactory receptor 4K14 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 4K14 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 4K14

Gene Name OR4K14

Cellular localization Cell membrane; Multi-pass membrane protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 37kD


Human Gene ID 122740

Human Swiss-Prot Number Q8NGD5

Alternative Names OR4K14; Olfactory receptor 4K14; Olfactory receptor OR14-22

Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using Olfactory receptor 4K14 Polyclonal Antibody diluted at 1:2000






Explore. Bioreagents.



Western blot analysis of lysates from COS7 and HUVEC cells, using OR4K14 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR4K14 antibody.