


p35 rabbit pAb**Cat#: orb765985 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	p35 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human CDK5R1. AA range:11-60
Specificity	p35 Polyclonal Antibody detects endogenous levels of p35 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cyclin-dependent kinase 5 activator 1
Gene Name	CDK5R1
Cellular localization	[Cyclin-dependent kinase 5 activator 1, p35]: Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cell projection, neuron projection . In the primary cortical neurons, p35 is present in the peripheries and nerve terminals. .; [Cyclin-dependent kinase 5 activator 1, p25]: Nucleus . Cytoplasm, perinuclear region . Perikaryon . The conversion of p35 to p25 relocalizes the protein from the cell periphery to the cytoplasm, in nuclear and perinuclear regions (PubMed:18507738). In the primary cortical neurons, p25 is primarily concentrated in the cell soma and is largely absent from neurites (PubMed:18507738)..


Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	38kD
Human Gene ID	8851
Human Swiss-Prot Number	Q15078
Alternative Names	CDK5R1; CDK5R; NCK5A; Cyclin-dependent kinase 5 activator 1; CDK5 activator 1; Cyclin-dependent kinase 5 regulatory subunit 1; TPKII regulatory subunit
Background	The protein encoded by this gene (p35) is a neuron-specific activator of cyclin-dependent kinase 5 (CDK5); the activation of CDK5 is required for proper development of the central nervous system. The p35 form of this protein is proteolytically cleaved by calpain, generating a p25 form. The cleavage of p35 into p25 results in relocalization of the protein from the cell periphery to nuclear and perinuclear regions. P25 deregulates CDK5 activity by prolonging its activation and changing its cellular location. The p25 form accumulates in the brain neurons of patients with Alzheimer's disease. This accumulation correlates with an increase in CDK5 kinase activity, and may lead to aberrantly phosphorylated forms of the microtubule-associated protein tau, which contributes to Alzheimer's disease. [provided by RefSeq, Jul 2008],




Immunofluorescence analysis of HeLa cell. 1, p35 Polyclonal Antibody (red) was diluted at 1:200 (4° overnight). Caspase 9 Monoclonal Antibody (3-20) (green) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog: RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog: RS3208 was diluted at 1:1000 (room temperature, 50min).



Western Blot analysis of various cells using p35 Polyclonal Antibody diluted at 1:1000



Western Blot analysis of HELA cells using p35 Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of HeLa cells, using CDK5R1 Antibody. The picture on the right is blocked with the synthesized peptide.