

Ribosomal Protein S6 rabbit pAb**Cat#: orb766252 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	Ribosomal Protein S6 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human S6 Ribosomal Protein. AA range:191-240
Specificity	Ribosomal Protein S6 Polyclonal Antibody detects endogenous levels of Ribosomal Protein S6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	40S ribosomal protein S6
Gene Name	RPS6
Cellular localization	nucleus,nucleoplasm,nucleolus,cytoplasm,cytosol,ribosome,polysome,small ribosomal subunit,membrane,cytosolic small ribosomal subunit,dendrite,intracellular ribonucleoprotein complex,cytoplasmic ribonucleoprotein granu
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	28kD
Human Gene ID	6194
Human Swiss-Prot Number	P62753
Alternative Names	RPS6; OK/SW-cl.2; 40S ribosomal protein S6; Phosphoprotein NP33

Background


Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed




Western Blot analysis of various cells using Ribosomal Protein S6 Polyclonal Antibody diluted at 1:2000



Western Blot analysis of HeLa cells using Ribosomal Protein S6 Polyclonal Antibody diluted at 1:2000



Western Blot analysis of customer's (cat sample) using Ribosomal Protein S6 Polyclonal Antibody diluted at 1:2000



The picture was kindly provided by our customer

Rps6