



RIN3 rabbit pAb

Cat#: orb766256 (Manual)

For research use only. Not intended for diagnostic use.

Product Name RIN3 rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000

Immunogen The antiserum was produced against synthesized peptide derived from

human RIN3. AA range:391-440

Specificity RIN3 Polyclonal Antibody detects endogenous levels of RIN3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Ras and Rab interactor 3

Gene Name RIN3

Cellular localization Cytoplasmic vesicle . Early endosome . Activation of tyrosine

phosphorylation signaling induces translocation to cytoplasmic vesicles. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 107kD


Human Gene ID 79890

Human Swiss-Prot Number Q8TB24


Alternative Names RIN3; Ras and Rab interactor 3; Ras interaction/interference protein 3

Summary: This protein encoded by this gene is a member of the RIN family **Background**

of Ras interaction-interference proteins, which are binding partners to the RAB5 small GTPases. The protein functions as a guanine nucleotide exchange for RAB5B and RAB31. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016],




Western Blot analysis of various cells using RIN3 Polyclonal Antibody



Western blot analysis of lysates from HUVEC and K562 cells, using RIN3 Antibody. The lane on the right is blocked with the synthesized peptide.







Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at $1:200(4^{\circ} \text{ overnight})$. 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).