



## TNF-α rabbit pAb

**Cat#: orb766498 (Manual)** 

For research use only. Not intended for diagnostic use.

**Product Name** TNF-α rabbit pAb

**Host species** Rabbit

**Applications** WB;IHC;IF;ELISA

**Species Cross-Reactivity** Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. IF 1:100-

300 Not yet tested in other applications.

The antiserum was produced against synthesized peptide derived from **Immunogen** 

human TNFA. AA range:141-190

TNF-α Polyclonal Antibody detects endogenous levels of TNF-α protein. **Specificity** 

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** 

Tumor necrosis factor (Cachectin) (TNF-alpha) (Tumor necrosis factor ligand superfamily member 2) (TNF-a) [Cleaved into: Tumor necrosis factor, membrane form (N-terminal fragment) (NTF); Intracellular **Protein Name** 

TNF TNFA TNFSF2 Gene Name

Cellular localization Cell membrane; Single-pass type II membrane protein.; [Tumor necrosis

factor, membrane form]: Membrane; Single-pass type II membrane protein.; [Tumor necrosis factor, soluble form]: Secreted .; [C-domain 1]: Secreted.;

[C-domain 2]: Secreted.





**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal

Concentration 1 mg/ml

Observed band 16kD

Human Gene ID 7124

**Human Swiss-Prot Number** P01375


Alternative Names TNF; TNFA; TNFSF2; Tumor necrosis factor; Cachectin; TNF-alpha;

Tumor necrosis factor ligand superfamily member 2; TNF-a

Background This gene encodes a multifunctional proinflammatory cytokine that belongs

to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.

[provided by RefSeq, Jul 2008],



Immunofluorescence analysis of A549. 1,primary Antibody(red) was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.