

β-1,4-Gal-T5 rabbit pAb

Cat#: orb766625 (Manual)


For research use only. Not intended for diagnostic use.

Product Name	β-1,4-Gal-T5 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human B4GALT5. AA range:321-370
Specificity	β-1,4-Gal-T5 Polyclonal Antibody detects endogenous levels of β-1,4-Gal-T5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Beta-1,4-galactosyltransferase 5
Gene Name	B4GALT5
Cellular localization	Golgi apparatus, Golgi stack membrane ; Single-pass type II membrane protein. Golgi apparatus . Trans cisternae of Golgi stack. .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal


Concentration	1 mg/ml
Observed band	40kD
Human Gene ID	9334
Human Swiss-Prot Number	O43286
Alternative Names	B4GALT5; Beta-1; 4-galactosyltransferase 5; Beta-1,4-GalTase 5; Beta4Gal-T5; b4Gal-T5; Beta-1,4-GalT II; UDP-Gal:beta-GlcNAc beta-1,4-galactosyltransferase 5; UDP-galactose:beta-N-acetylglucosamine beta-1,4-galactosyltransferase 5

Background

This gene is one of seven beta-1,4-galactosyltransferase (beta4GalT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. The function of the enzyme encoded by this gene is not clear. This gene was previously designated as B4GALT4 but was renamed



Western Blot analysis of various cells using β -1,4-Gal-T5 Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western Blot analysis of 293 cells using β -1,4-Gal-T5 Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000