

ASIC3 rabbit pAb**Cat#: orb766805 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	ASIC3 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human ASIC3. AA range:191-240
Specificity	ASIC3 Polyclonal Antibody detects endogenous levels of ASIC3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Acid-sensing ion channel 3
Gene Name	ASIC3
Cellular localization	Cell membrane ; Multi-pass membrane protein . Cytoplasm . Cell surface expression may be stabilized by interaction with LIN7B and cytoplasmic retention by interaction with DLG4. In part cytoplasmic in cochlea cells (By similarity). .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	58kD
Human Gene ID	9311
Human Swiss-Prot Number	Q9UHC3
Alternative Names	ASIC3; ACCN3; SLNAC1; TNAC1; Acid-sensing ion channel 3; ASIC3; hASIC3; Amiloride-sensitive cation channel 3; Neuronal amiloride-sensitive cation channel 3; Testis sodium channel 1; hTNaC1

Background

This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2012],



Western Blot analysis of HepG2 cells using ASIC3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from HepG2 cells, using ASIC3 Antibody.