



## CD32-B/C rabbit pAb

Cat#: orb766851 (Manual)

For research use only. Not intended for diagnostic use.

Product Name CD32-B/C rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from the

Internal region of human FCGR2B/C. AA range: 141-190

Specificity CD32-B/C Polyclonal Antibody detects endogenous levels of CD32-B/C

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Low affinity immunoglobulin gamma Fc region receptor II-b/Low affinity

immunoglobulin gamma Fc region receptor II-c

Gene Name FCGR2B/FCGR2C

Cellular localization Cell membrane; Single-pass type I membrane protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





1 mg/ml Concentration

**Observed band** 35kD

**Human Gene ID** 2213

P31994 **Human Swiss-Prot Number** 


**Alternative Names** 

FCGR2B; CD32; FCG2; IGFR2; Low affinity immunoglobulin gamma Fc region receptor II-b; IgG Fc receptor II-b; CDw32; Fc-gamma RII-b; Fc-gamma-RIIb; FcRII-b; CD32; FCGR2C; CD32; FCG2; IGFR2;Low affinity immunoglobulin gamma Fc region receptor II-c; IgG Fc re

Low affinity immunoglobulin gamma Fc region receptor II-b/Low affinity immunoglobulin gamma Fc region receptor II-c encoded by FCGR2B is a **Background** 

low affinity receptor for the Fc region of immunoglobulin gamma complexes. The encoded protein is involved in the phagocytosis of immune complexes and in the regulation of antibody production by B-cells. Variations in this gene may increase susceptibility to systemic lupus erythematosus (SLE). Several transcript variants encoding different isoforms

have been found for this gene.



Western Blot analysis of K562 cells using CD32-B/C Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000