



SPARC rabbit pAb

Cat#: orb767126 (Manual)

For research use only. Not intended for diagnostic use.

Product Name SPARC rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from the

Internal region of human SPARC. AA range: 141-190

Specificity SPARC Polyclonal Antibody detects endogenous levels of SPARC protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name SPARC

Gene Name SPARC

Cellular localization Secreted, extracellular space, extracellular matrix, basement membrane . In

or around the basement membrane.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 34kD

Human Gene ID 6678

Human Swiss-Prot Number P09486


Alternative Names SPARC; ON; SPARC; Basement-membrane protein 40; BM-40;

Osteonectin; ON; Secreted protein acidic and rich in cysteine


Background This gene encodes a cysteine-rich acidic matrix-associated protein. The

encoded protein is required for the collagen in bone to become calcified but is also involved in extracellular matrix synthesis and promotion of changes to cell shape. The gene product has been associated with tumor suppression but has also been correlated with metastasis based on changes to cell shape which can promote tumor cell invasion. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun

2015],



Western Blot analysis of HeLa, HepG2 cells using SPARC Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-muscle, antibody was diluted at 1:100



