



CD85d rabbit pAb

Cat#: orb767200 (Manual)

For research use only. Not intended for diagnostic use.

Product Name CD85d rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from the

Internal region of human LILRB2. AA range:121-170

Specificity CD85d Polyclonal Antibody detects endogenous levels of CD85d protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Leukocyte immunoglobulin-like receptor subfamily B member 2

Gene Name LILRB2

Cellular localization Cell membrane ; Single-pass type I membrane protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 66kD

Human Gene ID 10288

Human Swiss-Prot Number Q8N423

Alternative Names

LILRB2; ILT4; LIR2; MIR10; Leukocyte immunoglobulin-like receptor subfamily B member 2; LIR-2; Leukocyte immunoglobulin-like receptor 2;

subfamily B member 2; LIR-2; Leukocyte immunoglobulin-like receptor 2; CD85 antigen-like family member D; Immunoglobulin-like transcript 4; ILT-


4; Monocyte/macrophage immunoglobulin-like re

Background This gene is a member of the leukocyte immunoglobulin-like receptor (LIR)

family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the

immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008],



Western Blot analysis of Hela cells using CD85d Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000