



## IL-10Rα rabbit pAb

Cat#: orb767235 (Manual)

For research use only. Not intended for diagnostic use.

**Product Name** IL-10Rα rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from the

Internal region of human IL10RA. AA range:151-200

Specificity IL-10Rα Polyclonal Antibody detects endogenous levels of IL-10Rα protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interleukin-10 receptor subunit alpha

Gene Name IL10RA

Cellular localization Cell membrane ; Single-pass type I membrane protein. Cytoplasm .

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





1 mg/ml

**Observed band** 

Concentration

63kD

**Human Gene ID** 

3587

**Human Swiss-Prot Number** 


Q13651

**Alternative Names** 

IL10RA; IL10R; Interleukin-10 receptor subunit alpha; IL-10 receptor subunit alpha; IL-10R subunit alpha; IL-10RA; CDw210a; Interleukin-10 receptor subunit 1; IL-10R subunit 1; IL-10R1; CD210

**Background** 

The protein encoded by this gene is a receptor for interleukin 10. This protein is structurally related to interferon receptors. It has been shown to mediate the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines. This receptor is reported to promote survival of progenitor myeloid cells through the insulin receptor substrate-2/PI 3-kinase/AKT pathway. Activation of this receptor leads to tyrosine phosphorylation of JAK1 and TYK2 kinases. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene. [provided by RefSeq, Jan 2009],



Western Blot analysis of PC12 cells using IL-10R $\alpha$  Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000