



CKR-2 rabbit pAb

Cat#: orb767269 (Manual)

For research use only. Not intended for diagnostic use.

Product Name CKR-2 rabbit pAb

Host species Rabbit

Applications WB;IF;ELISA

Species Cross-Reactivity Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:

1/5000. Not yet tested in other applications.

Immunogen Synthesized peptide derived from CKR-2 . at AA range: 250-330

Specificity CKR-2 Polyclonal Antibody detects endogenous levels of CKR-2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name C-C chemokine receptor type 2

Gene Name CCR2

Cellular localization

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

Concentration 1 mg/ml

Observed band 46kD


Human Gene ID

Human Swiss-Prot Number

CCR2; CMKBR2; C-C chemokine receptor type 2; C-C CKR-2; CC-CKR-2; CCR-2; CCR2; Monocyte chemoattractant protein 1 receptor; MCP-1-R; CD antigen CD192 **Alternative Names**

Background

CCR2 (C-C Motif Chemokine Receptor 2) is a Protein Coding gene. Diseases associated with CCR2 include idiopathic anterior uveitis and cd3zeta deficiency. Among its related pathways are Signaling by GPCR and A-beta Pathways: uptake and Degradation. GO annotations related to this gene include G-protein coupled receptor activity and chemokine receptor activity. An important paralog of this gene is CCR9. eceptor for the CCL2, CCL7 and CCL13 chemokines. Transduces a signal by increasing intracellular calcium ion levels. Alternative coreceptor with CD4 for HIV-1 infaction hard line CC receptor (CCR2) coreceptor with constitution and constitution of CC. infection. hemokine CC receptors (CCRs) predominantly recognize CC chemokines. CC chemokines are distinguished by having four conserved cysteines, with the first two cysteines being adjacent to each other. There are 10 chemokine CC receptors.



Western Blot analysis of K562 cells using CKR-2 Polyclonal Antibody