



B-ATF rabbit pAb

Cat#: orb767377 (Manual)

For research use only. Not intended for diagnostic use.

Product Name B-ATF rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

Recommended dilutions Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human BATF. AA range:10-59

Specificity B-ATF Polyclonal Antibody detects endogenous levels of B-ATF protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Basic leucine zipper transcriptional factor ATF-like

Gene Name BATF

Cellular localization Nucleus . Cytoplasm . Present in the nucleus and cytoplasm, but shows

increased nuclear translocation after activation of T-cells. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





1 mg/mlConcentration

Observed band


Human Gene ID 10538

Human Swiss-Prot Number Q16520


BATF; Basic leucine zipper transcriptional factor ATF-like; B-cell-activating transcription factor; B-ATF; SF-HT-activated gene 2 protein; SFA-2 **Alternative Names**

The protein encoded by this gene is a nuclear basic leucine zipper protein that belongs to the AP-1/ATF superfamily of transcription factors. The **Background**

leucine zipper of this protein mediates dimerization with members of the Jun family of proteins. This protein is thought to be a negative regulator of AP-1/ATF transcriptional events. [provided by RefSeq, Jul 2008],



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absor



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using BATF Antibody. The picture on the right is blocked with the synthesized peptide.