



CA VB rabbit pAb

Cat#: orb767498 (Manual)


For research use only. Not intended for diagnostic use.

Product Name	CA VB rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human CA5B. AA range:241-290
Specificity	CA VB Polyclonal Antibody detects endogenous levels of CA VB protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
Formulation Storage	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Store at -20°C. Avoid repeated freeze-thaw cycles.
	azide
Storage	azide Store at -20°C. Avoid repeated freeze-thaw cycles.
Storage Protein Name	azide Store at -20°C. Avoid repeated freeze-thaw cycles. Carbonic anhydrase 5B mitochondrial
Storage Protein Name Gene Name	azide Store at -20°C. Avoid repeated freeze-thaw cycles. Carbonic anhydrase 5B mitochondrial CA5B




www.biorbyt.com

Concentration	1 mg/ml
Observed band	38kD
Human Gene ID	11238
Human Swiss-Prot Number	Q9Y2D0
Alternative Names	CA5B; Carbonic anhydrase 5B; mitochondrial; Carbonate dehydratase VB; Carbonic anhydrase VB; CA-VB
Background	Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. CA VB is localized in the mitochondria and shows the highest sequence similarity to the other mitochondrial CA, CA VA. It has a wider tissue distribution than CA VA, which is restricted to the liver. The differences in tissue distribution suggest that the two mitochondrial carbonic anhydrases evolved to assume different physiologic roles. [provided by RefSeq, Jul 2008],







www.biorbyt.com



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Immunofluorescence analysis of MCF7 cells, using CA5B Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, using CA5B Antibody. The lane on the right is blocked with the synthesized peptide.