



## CKR-4 rabbit pAb


## Cat#: orb767582 (Manual)

For research use only. Not intended for diagnostic use.

Product Name	CKR-4 rabbit pAb
Host species	Rabbit
Applications	IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human CCR4. AA range:211-260
Specificity	CKR-4 Polyclonal Antibody detects endogenous levels of CKR-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
Formulation Storage	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Store at -20°C. Avoid repeated freeze-thaw cycles.
	azide
Storage	azide Store at -20°C. Avoid repeated freeze-thaw cycles.
Storage Protein Name	azide Store at -20°C. Avoid repeated freeze-thaw cycles. C-C chemokine receptor type 4
Storage Protein Name Gene Name	azide Store at -20°C. Avoid repeated freeze-thaw cycles. C-C chemokine receptor type 4 CCR4



Concentration	1 mg/ml
Observed band	41kD
Human Gene ID	1233
Human Swiss-Prot Number	P51679
Alternative Names	CCR4; CMKBR4; C-C chemokine receptor type 4; C-C CKR-4; CC-CKR-4; CCR-4; CCR-4; CCR4; K5-5; CD antigen CD194
Background	The protein encoded by this gene belongs to the G-protein-coupled receptor family. It is a receptor for the CC chemokine - MIP-1, RANTES, TARC and MCP-1. Chemokines are a group of small polypeptide, structurally related molecules that regulate cell trafficking of various types of leukocytes. The chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. [provided by RefSeq, Jul 2008],



Immunofluorescence analysis of HUVEC cells, using CCR4 Antibody. The picture on the right is blocked with the synthesized peptide.