



ACOT12 rabbit pAb

Cat#: orb767650 (Manual)

For research use only. Not intended for diagnostic use.

Product Name ACOT12 rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000.

ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human ACOT12. AA range:281-330

Specificity ACOT12 Polyclonal Antibody detects endogenous levels of ACOT12

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Acyl-coenzyme A thioesterase 12

Gene Name ACOT12

Cellular localization Cytoplasm, cytosol.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

1 mg/mlConcentration

Observed band

Human Gene ID 134526

Human Swiss-Prot Number Q8WYK0


Alternative Names ACOT12; CACH; CACH1; STARD15; Acyl-coenzyme A thioesterase 12;

Acyl-CoA thioesterase 12; Acyl-CoA thioester hydrolase 12; Cytoplasmic acetyl-CoA hydrolase 1; CACH-1; hCACH-1; START domain-containing


protein 15; StARD15

Background

catalytic activity: Acetyl-CoA + H(2)O = CoA + acetate., function: Hydrolyzes acetyl-CoA to acetate and CoA., pathway: Carbohydrate metabolism; pyruvate metabolism., similarity: Contains 1 START domain., similarity: Contains 2 acyl coenzyme A hydrolase domains., subunit: Homodimer or homotetramer.,



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbe




Immunofluorescence analysis of A549 cells, using ACOT12 Antibody. The picture on the right is blocked with the synthesized peptide.





Explore. Bioreagents.



Immunohistochemistry analysis of paraffin-embedded human liver carcinoma tissue, using ACOT12 Antibody. The picture on the right is blocked with the synthesized peptide.