

PARP-4 rabbit pAb

Cat#: orb767710 (Manual)


For research use only. Not intended for diagnostic use.

Product Name	PARP-4 rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human PARP4. AA range:1151-1200
Specificity	PARP-4 Polyclonal Antibody detects endogenous levels of PARP-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Poly [ADP-ribose] polymerase 4
Gene Name	PARP4
Cellular localization	Cytoplasm . Nucleus . Cytoplasm, cytoskeleton, spindle . Also found in the nucleus, associated with mitotic spindles. .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal


Concentration	1 mg/ml
Observed band	
Human Gene ID	143
Human Swiss-Prot Number	Q9UKK3
Alternative Names	PARP4; ADPRTL1; KIAA0177; PARPL; Poly [ADP-ribose] polymerase 4; PARP-4; 193 kDa vault protein; ADP-ribosyltransferase diphtheria toxin-like 4; ARTD4; PARP-related/Alphal-related H5/proline-rich; PH5P; Vault poly(ADP-ribose) polymerase; VP

Background

This gene encodes poly(ADP-ribosyl)transferase-like 1 protein, which is capable of catalyzing a poly(ADP-ribosyl)ation reaction. This protein has a catalytic domain which is homologous to that of poly (ADP-ribosyl) transferase, but lacks an N-terminal DNA binding domain which activates the C-terminal catalytic domain of poly (ADP-ribosyl) transferase. Since this protein is not capable of binding DNA directly, its transferase activity may be activated by other factors such as protein-protein interaction mediated by the extensive carboxyl terminus. [provided by RefSeq, Jul 2008],



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using PARP4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using FAF1 antibody.



Explore. Bioreagents.

www.biorbyt.com