



Olfactory receptor 10A5 rabbit pAb

Cat#: orb767726 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Olfactory receptor 10A5 rabbit pAb

Host species Rabbit

Applications IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human OR10A5. AA range:200-249

Specificity Olfactory receptor 10A5 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 10A5 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 10A5

Gene Name OR10A5

Cellular localization Cell membrane; Multi-pass membrane protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





1 mg/mlConcentration

Observed band

Human Gene ID 144124


Human Swiss-Prot Number Q9H207

OR10A5; OR10A1; Olfactory receptor 10A5; HP3; Olfactory receptor 10A1; Olfactory receptor 11-403; OR11-403; Olfactory receptor-like protein **Alternative Names**

JCG6

Background Olfactory receptors interact with odorant molecules in the nose, to initiate a

neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Immunofluorescence analysis of MCF7 cells, using OR10A5 Antibody. The picture on the right is blocked with the synthesized peptide.