

eIF4G (phospho Ser1148) rabbit pAb

Cat#: orb767959 (Manual)


For research use only. Not intended for diagnostic use.

Product Name	eIF4G (phospho Ser1148) rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000 ,Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human eIF4G around the phosphorylation site of Ser1108. AA range:1074-1123
Specificity	Phospho-eIF4G (S1148) Polyclonal Antibody detects endogenous levels of eIF4G protein only when phosphorylated at S1148.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Eukaryotic translation initiation factor 4 gamma 1
Gene Name	EIF4G1
Cellular localization	Cytoplasm, Stress granule .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal


Concentration	1 mg/ml
Observed band	
Human Gene ID	1981
Human Swiss-Prot Number	Q04637
Alternative Names	EIF4G1; EIF4F; EIF4G; EIF4GI; Eukaryotic translation initiation factor 4 gamma 1; eIF-4-gamma 1; eIF-4G 1; eIF-4G1; p220

Background

The protein encoded by this gene is a component of the multi-subunit protein complex EIF4F. This complex facilitates the recruitment of mRNA to the ribosome, which is a rate-limiting step during the initiation phase of protein synthesis. The recognition of the mRNA cap and the ATP-dependent unwinding of 5'-terminal secondary structure is catalyzed by factors in this complex. The subunit encoded by this gene is a large scaffolding protein that contains binding sites for other members of the EIF4F complex. A domain at its N-terminus can also interact with the poly(A)-binding protein, which may mediate the circularization of mRNA during translation. Alternative splicing results in multiple transcript variants, some of which are derived from alternative promoter usage. [provided by RefSeq, Aug 2010],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using eIF4G (Phospho-Ser1108) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using eIF4G (Phospho-Ser1108) Antibody. The picture on the right is blocked with the phospho peptide.