



PI 3-kinase p101 rabbit pAb

Cat#: orb768229 (Manual)

For research use only. Not intended for diagnostic use.

Product Name PI 3-kinase p101 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human PIK3R5. AA range:695-744

Specificity PI 3-kinase p101 Polyclonal Antibody detects endogenous levels of PI 3-

kinase p101 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Phosphoinositide 3-kinase regulatory subunit 5

Gene Name PIK3R5

Cellular localization Nucleus . Cytoplasm . Cell membrane ; Peripheral membrane protein .

Predominantly localized in the nucleus in absence of PIK3CG/p120.

Colocalizes with PIK3CG/p120 in the cytoplasm. Translocated to the plasma

membrane in a beta-gamma G protein-dependent manner. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.





Clonality Polyclonal

Concentration 1 mg/ml

Observed band 100kD

Human Gene ID 23533

Human Swiss-Prot Number Q8WYR1

Alternative Names PIK3R5; Phosphoinositide 3-kinase regulatory subunit 5; PI3-kinase


regulatory subunit 5; PI3-kinase p101 subunit; Phosphatidylinositol 4; 5-bisphosphate 3-kinase regulatory subunit; PtdIns-3-kinase regulatory subunit;

Protein FOAP-2; PtdIns-

Background Phosphatidylinositol 3-kinases (PI3Ks) phosphorylate the inositol ring of

phosphatidylinositol at the 3-prime position, and play important roles in cell growth, proliferation, differentiation, motility, survival and intracellular trafficking. The PI3Ks are divided into three classes: I, II and III, and only the class I PI3Ks are involved in oncogenesis. This gene encodes the 101 kD regulatory subunit of the class I PI3K gamma complex, which is a dimeric enzyme, consisting of a 110 kD catalytic subunit gamma and a regulatory subunit of either 55, 87 or 101 kD. This protein recruits the catalytic subunit from the cytosol to the plasma membrane through high-affinity interaction with G-beta-gamma proteins. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been found. [provided by

RefSeq, Oct 2011],



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using PIK3R5 Antibody. The picture on the right is blocked with the synthesized peptide.