



HMG-17 (phospho Ser29) rabbit pAb

Cat#: orb768620 (Manual)

For research use only. Not intended for diagnostic use.

Product Name	HMG-17 (phospho Ser29) rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human HMG17 around the phosphorylation site of Ser29. AA range:1-50
Specificity	Phospho-HMG-17 (S29) Polyclonal Antibody detects endogenous levels of HMG-17 protein only when phosphorylated at S29.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Non-histone chromosomal protein HMG-17
Gene Name	HMGN2
Cellular localization	Nucleus . Cytoplasm . Cytoplasmic enrichment upon phosphorylation.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Clonality	Polyclonal



www.biorbyt.com

Concentration	1 mg/ml
Observed band	
Human Gene ID	3151
Human Swiss-Prot Number	P05204
Alternative Names	HMGN2; HMG17; Non-histone chromosomal protein HMG-17; High mobility group nucleosome-binding domain-containing protein 2
Background	high mobility group nucleosomal binding domain 2(HMGN2) Homo sapiens The protein encoded by this gene binds nucleosomal DNA and is associated with transcriptionally active chromatin. Along with a similar protein, HMGN1, the encoded protein may help maintain an open chromatin configuration around transcribable genes. The protein has also been found to have antimicrobial activity against bacteria, viruses and fungi. [provided by RefSeq, Oct 2014],



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by i