



## Olfactory receptor 51B6 rabbit pAb

**Cat#: orb768922 (Manual)** 

For research use only. Not intended for diagnostic use.

Product Name Olfactory receptor 51B6 rabbit pAb

Host species Rabbit

Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:

1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human OR51B6. AA range:66-115

Specificity Olfactory receptor 51B6 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 51B6 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 51B6

Gene Name OR51B6

Cellular localization Cell membrane; Multi-pass membrane protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Concentration 1 mg/ml

Observed band 35kD


Human Gene ID 390058

Human Swiss-Prot Number Q9H340

Alternative Names OR51B6; Olfactory receptor 51B6; Odorant receptor HOR5'beta6

**Background** 

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Immunofluorescence analysis of MCF7 cells, using OR51B6 Antibody. The picture on the right is blocked with the synthesized peptide.





Explore. Bioreagents.



Western blot analysis of lysates from K562 cells, using OR51B6 Antibody. The lane on the right is blocked with the synthesized peptide.