

Olfactory receptor 51I2 rabbit pAb

Cat#: orb768925 (Manual)


For research use only. Not intended for diagnostic use.

| | |
|---------------------------------|--|
| Product Name | Olfactory receptor 51I2 rabbit pAb |
| Host species | Rabbit |
| Applications | WB;IF;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. |
| Immunogen | The antiserum was produced against synthesized peptide derived from human OR51I2. AA range:201-250 |
| Specificity | Olfactory receptor 51I2 Polyclonal Antibody detects endogenous levels of Olfactory receptor 51I2 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Olfactory receptor 51I2 |
| Gene Name | OR51I2 |
| Cellular localization | Cell membrane; Multi-pass membrane protein. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Clonality | Polyclonal |


| | |
|--------------------------------|---|
| Concentration | 1 mg/ml |
| Observed band | 35kD |
| Human Gene ID | 390064 |
| Human Swiss-Prot Number | Q9H344 |
| Alternative Names | OR51I2; Olfactory receptor 51I2; Odorant receptor HOR5"beta12; Olfactory receptor OR11-38 |

Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from HUVEC cells, using OR51I2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR51I2 antibody.