



Laminin α-5 rabbit pAb

Cat#: orb768952 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Laminin α-5 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. IF 1:100-

300 Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human LAMA5. AA range:2381-2430

Specificity Laminin α -5 Polyclonal Antibody detects endogenous levels of Laminin α -5

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Laminin subunit alpha-5

Gene Name LAMA5

Cellular localization Secreted, extracellular space, extracellular matrix, basement membrane.

Major component.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration 1 mg/ml

Observed band 400kD

Human Gene ID 3911


Human Swiss-Prot Number O15230

Alternative Names LAMA5; KIAA0533; KIAA1907; Laminin subunit alpha-5; Laminin-10

subunit alpha; Laminin-11 subunit alpha; Laminin-15 subunit alpha

Background

This gene encodes one of the vertebrate laminin alpha chains. Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins are composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively) and they form a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. The protein encoded by this gene is the alpha-5 subunit of of laminin-10 (laminin-511), laminin-11 (laminin-521) and laminin-15 (laminin-523). [provided by RefSeq, Jun 2013],




Immunofluorescence analysis of A549. 1,primary Antibody was diluted at $1:200(4^{\circ}\text{C}\ \text{overnight})$. 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at $1:1000(\text{room\ temperature},50\text{min})$.3, Picture B: DAPI(blue) 10min.







Explore. Bioreagents.



Western Blot analysis of RAW HepG2 A549 cells using Laminin α -5 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using LAMA5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, $4^{\circ} over\ night$