

**LIMK-2 (phospho Ser283) rabbit pAb****Cat#: orb768990 (Manual)**


For research use only. Not intended for diagnostic use.

<b>Product Name</b>	LIMK-2 (phospho Ser283) rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat;Monkey
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human LIMK2 around the phosphorylation site of Ser283. AA range:249-298
<b>Specificity</b>	Phospho-LIMK-2 (S283) Polyclonal Antibody detects endogenous levels of LIMK-2 protein only when phosphorylated at S283.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	LIM domain kinase 2
<b>Gene Name</b>	LIMK2
<b>Cellular localization</b>	Cytoplasm, cytoskeleton, spindle . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .; [Isoform LIMK2a]: Cytoplasm . Nucleus .; [Isoform LIMK2b]: Cytoplasm . Cytoplasm, perinuclear region . Nucleus . Mainly present in the cytoplasm and is scarcely translocated to the nucleus. .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	72kD
<b>Human Gene ID</b>	3985
<b>Human Swiss-Prot Number</b>	P53671
<b>Alternative Names</b>	LIMK2; LIM domain kinase 2; LIMK-2

## Background


There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],



**Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using LIMK2 (Phospho-Ser283) Antibody**



Immunohistochemistry analysis of paraffin-embedded human brain, using LIMK2 (Phospho-Ser283) Antibody. The picture on the right is blocked with the phosphopeptide.



Western blot analysis of LIMK2 (Phospho-Ser283) Antibody. The lane on the right is blocked with the LIMK2 (Phospho-Ser283) peptide.