



## PAKα/β/γ rabbit pAb

**Cat#: orb769339 (Manual)** 

For research use only. Not intended for diagnostic use.

**Product Name** PAK $\alpha/\beta/\gamma$  rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

**Recommended dilutions** Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human PAK1/2/3. ÅA range:111-160

Specificity PAK $\alpha/\beta/\gamma$  Polyclonal Antibody detects endogenous levels of PAK $\alpha/\beta/\gamma$ 

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/threonine-protein kinase PAK 1

Gene Name PAK1/PAK2/PAK3

Cellular localization Cytoplasm . Cell junction, focal adhesion . Cell projection, lamellipodium .

Cell membrane . Cell projection, ruffle membrane . Cell projection, invadopodium . Nucleus, nucleoplasm . Chromosome . Cytoplasm,

cytoskeleton, microtubule organizing center, centrosome. Colocalizes with RUFY3, F-actin and other core migration components in invadopodia at the cell periphery (PubMed:25766321). Recruited to the cell membrane by interaction with CDC42 and RAC1. Recruited to focal adhesions upon activation. Colocalized with CIB1 within membrane ruffles during cell spreading upon readhesion to fibronectin. Upon DNA damage, translocates to the nucleoplasm when phosphorylated at Thr-212 where is co-recruited with MORC2 on damaged chromatin (PubMed:23260667). Localization to



## www.biorbyt.com

the centrosome does not depen

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

epitope-specific immunogen. chromatography using

Polyclonal **Clonality** 

Concentration 1 mg/ml

**Observed band** 

**Human Gene ID** 5058/5062/5063

**Human Swiss-Prot Number** Q13153/Q13177/O75914


**Alternative Names** 

PAK1; Serine/threonine-protein kinase PAK 1; Alpha-PAK; p21-activated kinase 1; PAK-1; p65-PAK; PAK2; Serine/threonine-protein kinase PAK 2; Gamma-PAK; PAK65; S6/H4 kinase; p21-activated kinase 2; PAK-2; p58;

PAK3; OPHN3; Serine/threonine-p

**Background** 

This gene encodes a family member of serine/threonine p21-activating kinases, known as PAK proteins. These proteins are critical effectors that link RhoGTPases to cytoskeleton reorganization and nuclear signaling, and they serve as targets for the small GTP binding proteins Cdc42 and Rac. This specific family member regulates cell motility and morphology. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2010],



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using PAK1/2/3 Antibody. The picture on the right is blocked with the synthesized peptide.