



## CLLD7 rabbit pAb

**Cat#: orb769591 (Manual)** 

For research use only. Not intended for diagnostic use.

Product Name CLLD7 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human RCBTB1. AA range:251-300

Specificity CLLD7 Polyclonal Antibody detects endogenous levels of CLLD7 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name RCC1 and BTB domain-containing protein 1

Gene Name RCBTB1

Cellular localization Nucleus.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Concentration 1 mg/ml

**Observed band** 58kD

**Human Gene ID** 55213


**Human Swiss-Prot Number** Q8NDN9

**Alternative Names** RCBTB1; CLLD7; E4.5; RCC1 and BTB domain-containing protein 1;


Chronic lymphocytic leukemia deletion region gene 7 protein; CLL deletion region gene 7 protein; Regulator of chromosome condensation and BTB domain-containing protein 1

This gene encodes a protein with an N-terminal RCC1 domain and a C-terminal BTB (broad complex, tramtrack and bric-a-brac) domain. In rat, Background

over-expression of this gene in vascular smooth muscle cells induced cellular hypertrophy. In rat, the C-terminus of RCBTB1 interacts with the angiotensin II receptor-1A. In humans, this gene maps to a region of chromosome 13q that is frequently deleted in B-cell chronic lymphocytic leukemia and other lymphoid molicensories. [Provided by Baffes Lt 2000] leukemia and other lymphoid malignancies. [provided by RefSeq, Jul 2008],



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using RCBTB1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from LOVO cells, using RCBTB1 Antibody. The lane on the right is blocked with the synthesized peptide.



