



## Krs-1/2 rabbit pAb

Cat#: orb770180 (Manual)

For research use only. Not intended for diagnostic use.

**Product Name** Krs-1/2 rabbit pAb

**Host species** Rabbit

**Applications** WB;IHC;IF;ELISA

**Species Cross-Reactivity** Human; Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/5000. Not yet tested in other applications.

**Immunogen** The antiserum was produced against synthesized peptide derived from

human Mst1/2. AA range:149-198

Krs-1/2 Polyclonal Antibody detects endogenous levels of Krs-1/2 protein. **Specificity** 

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** 

**Protein Name** Serine/threonine-protein kinase 3/4

Gene Name STK3/STK4

Cytoplasm . Nucleus . The caspase-cleaved form cycles between nucleus and cytoplasm (PubMed:19525978, PubMed:11278283). Phosphorylation at Thr-117 leads to inhibition of nuclear translocation (PubMed:19525978). . Cellular localization

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.





Polyclonal **Clonality** 

Concentration 1 mg/ml

**Observed band** 56kD

**Human Gene ID** 6788/6789

**Human Swiss-Prot Number** Q13188/Q13043


**Alternative Names** 

STK3; KRS1; MST2; Serine/threonine-protein kinase 3; Mammalian STE20-like protein kinase 2; MST-2; STE20-like kinase MST2; Serine/threonine-protein kinase Krs-1; STK4; KRS2; MST1; Serine/threonine-protein kinase 4; Mammalian STE20-like prot

**Background** serine/threonine kinase 3(STK3) Homo sapiens This gene encodes a

serine/threonine protein kinase activated by proapoptotic molecules indicating the encoded protein functions as a growth suppressor. Cleavage of the protein product by caspase removes the inhibitory C-terminal portion. The N-terminal portion is transported to the nucleus where it homodimerizes to form the active kinase which promotes the condensation of chromatin during apoptosis. Multiple transcript variants encoding different isoforms

have been found for this gene. [provided by RefSeq, Jan 2012],



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Mst1/2 Antibody. The picture on the right is blocked with the synthesized peptide.





Explore. Bioreagents.

