



Actin-α/γ (phospho Tyr55/53) rabbit pAb

Cat#: orb770235 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Actin-α/γ (phospho Tyr55/53) rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human Actin-pan around the phosphorylation site of Tyr55/53. AA range:21-

70

Specificity Phospho-Actin-α/γ (Y55/53) Polyclonal Antibody detects endogenous levels

of Actin- α/γ protein only when phosphorylated at Y55/53.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Actin alpha cardiac muscle 1

Gene Name ACTC1

Cellular localization Cytoplasm, cytoskeleton.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





1 mg/mlConcentration

Observed band

Human Gene ID 70/71/72/58

Human Swiss-Prot Number P68032/P63261/P63267/P68133


Alternative Names ACTC1; ACTC; Actin; alpha cardiac muscle 1; Alpha-cardiac actin; ACTG1; ACTB; ACTG; Actin, cytoplasmic 2; Gamma-actin; ACTG2

ACTA3; ACTL3; ACTSG; Actin, gamma-enteric smooth muscle; Alpha-

actin-3; Gamma-2-actin; Smooth muscle gamma-actin;

Background

Actins are highly conserved proteins that are involved in various types of cell motility. Polymerization of globular actin (G-actin) leads to a structural filament (F-actin) in the form of a two-stranded helix. Each actin can bind to four others. The protein encoded by this gene belongs to the actin family which is comprised of three main groups of actin isoforms, alpha, beta, and gamma. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. Defects in this gene have been associated with idiopathic dilated cardiomyopathy (IDC) and familial hypertrophic cardiomyopathy (FHC). [provided by RefSeq, Jul 2008],



Immunohistochemistry analysis of paraffin-embedded human skeletal muscle, using Actin-pan (alpha/gamma) (Phospho-Tyr55/53) Antibody. The picture on the right is blocked with the phospho peptide.