



## EMR3 rabbit pAb

Cat#: orb770647 (Manual)

For research use only. Not intended for diagnostic use.

Product Name EMR3 rabbit pAb

Host species Rabbit

**Applications** WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:

1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human EMR3. AA range:603-652

Specificity EMR3 Polyclonal Antibody detects endogenous levels of EMR3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name EGF-like module-containing mucin-like hormone receptor-like 3

Gene Name EMR3

Cellular localization Cell membrane; Multi-pass membrane protein .; [Isoform 3]: Secreted.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Concentration 1 mg/ml

Observed band 70kD

Human Gene ID 84658


Human Swiss-Prot Number Q9BY15

Alternative Names EMR3; EGF-like module-containing mucin-like hormone receptor-like 3;

EGF-like module receptor 3

**Background** 

This gene encodes a member of the class B seven-span transmembrane (TM7) receptor family expressed predominantly by cells of the immune system. Family members are characterized by an extended extracellular region with a variable number of N-terminal epidermal growth factor (EGF)-like domains coupled to a TM7 domain via a mucin-like spacer domain. This gene is closely linked to the gene encoding egf-like molecule containing mucin-like hormone receptor 2 on chromosome 19. This protein may play a role in myeloid-myeloid interactions during immune and inflammatory responses. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2014],



Western blot analysis of lysates from Jurkat cells, using EMR3 Antibody. The lane on the right is blocked with the synthesized peptide.