

SNX4 rabbit pAb**Cat#: orb770741 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	SNX4 rabbit pAb
Host species	Rabbit
Applications	WB;IHC
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000;IHC-p 1:50-300
Immunogen	The antiserum was produced against synthesized peptide derived from human SNX4. AA range:191-240
Specificity	SNX4 Polyclonal Antibody detects endogenous levels of SNX4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Sorting nexin-4
Gene Name	SNX4
Cellular localization	Early endosome membrane ; Peripheral membrane protein ; Cytoplasmic side . Also detected on a juxtanuclear endocytic recycling compartment (ERC). .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	60kD
Human Gene ID	8723
Human Swiss-Prot Number	O95219
Alternative Names	SNX4; Sorting nexin-4

Background

This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein associated with the long isoform of the leptin receptor and with receptor tyrosine kinases for platelet-derived growth factor, insulin, and epidermal growth factor in cell cultures, but its function is unknown. This protein may form oligomeric complexes with family members. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene. [provided by RefSeq, Nov 2012],



Western blot analysis of the lysates from HeLa cells using SNX4 antibody.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).