



GABAB R2 rabbit pAb

Cat#: orb770938 (Manual)

For research use only. Not intended for diagnostic use.

Product Name GABAB R2 rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen Synthesized peptide derived from GABAB R2. at AA range: 830-910

GABAB R2 Polyclonal Antibody detects endogenous levels of GABAB R2 **Specificity**

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Protein Name Gamma-aminobutyric acid type B receptor subunit 2

Gene Name GABBR2

Cellular localization

Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein. Coexpression of GABBR1 and GABBR2 is required for GABBR1 maturation and transport

to the plasma membrane. In contrast, GABBR2 does not depend on

GABBR1 for transport to the cell membrane. .

The antibody was affinity-purified from rabbit antiserum by affinity-**Purification**

chromatography using epitope-specific immunogen.





Clonality Polyclonal

Concentration 1 mg/ml

Observed band 105kD

9568 **Human Gene ID**

Human Swiss-Prot Number O75899

GABBR2; GPR51; GPRC3B; Gamma-aminobutyric acid type B receptor subunit 2; GABA-B receptor 2; GABA-B-R2; GABA-BR2; GABABR2; Gb2; G-protein coupled receptor 51; HG20 **Alternative Names**

Background

The multi-pass membrane protein encoded by this gene belongs to the G-protein coupled receptor 3 family and GABA-B receptor subfamily. The GABA-B receptors inhibit neuronal activity through G protein-coupled second-messenger systems, which regulate the release of neurotransmitters, and the activity of ion channels and adenylyl cyclase. This receptor subunit forms an active heterodimeric complex with GABA-B receptor subunit 1, neither of which is effective on its own. Allelic variants of this gene have been associated with nicotine dependence. [provided by RefSeq, Jan 2010],