



hnRNP DL rabbit pAb

Cat#: orb771027 (Manual)

For research use only. Not intended for diagnostic use.

Product Name hnRNP DL rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human HNRPDL. ÂA range:241-290

Specificity hnRNP DL Polyclonal Antibody detects endogenous levels of hnRNP DL

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Heterogeneous nuclear ribonucleoprotein D-like

Gene Name HNRPDL

Cellular localization Nucleus . Cytoplasm . Shuttles between the nucleus and the cytoplasm in a

TNPO1-dependent manner. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





1 mg/mlConcentration

Observed band

Human Gene ID 9987

Human Swiss-Prot Number O14979


HNRPDL; JKTBP; Heterogeneous nuclear ribonucleoprotein D-like; hnRNP D-like; hnRNP DL; AU-rich element RNA-binding factor; JKT41-binding **Alternative Names**

protein; Protein laAUF1

heterogeneous nuclear ribonucleoprotein D like(HNRNPDL) Homo sapiens **Background**

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to In the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two RRM domains that bind to RNAs. Three alternatively spliced transcript variants have been described for this gene. One of the variants is probably not translated because the transcript is a candidate for nonsense mediated not translated because the transcript is a candidate for nonsense-mediated mRNA decay. The protein isoforms encoded by this gene are similar to its

family member



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using HNRPDL Antibody. The picture on the right is blocked with the synthesized peptide.