



## Vitronectin rabbit pAb

Cat#: orb771261 (Manual)

For research use only. Not intended for diagnostic use.

**Product Name** Vitronectin rabbit pAb

**Host species** Rabbit

**Applications** WB;ELISA

**Species Cross-Reactivity** Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

**Immunogen** The antiserum was produced against synthesized peptide derived from the

Internal region of human VTN. AA range:50-100

Vitronectin Polyclonal Antibody detects endogenous levels of Vitronectin **Specificity** 

protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** 

**Protein Name** Vitronectin

Gene Name VTN

Cellular localization

Secreted, extracellular space .; Parasitophorous vacuole . (Microbial infection) In P.falciparum-infected red blood cells, VTN internalization is detected at the early trophozoite stage (PubMed:29567995). Colocalizes with SERA5 at the scrizont stage and with SERA5 P47 at the merozoite surface

(PubMed:29567995). .

The antibody was affinity-purified from rabbit antiserum by affinity-**Purification** 

chromatography using epitope-specific immunogen.





**Clonality** Polyclonal

Concentration 1 mg/ml

**Observed band** 55kD


7448 **Human Gene ID** 

**Human Swiss-Prot Number** P04004

VTN; Vitronectin; VN; S-protein; Serum-spreading factor; V75 **Alternative Names** 

Background The protein encoded by this gene is a member of the pexin family. It is found

in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond. [provided by RefSeq, Jul 2008],



Western blot analysis of mouse-brain mouse-lung lysis using VTN antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000