



## CD88 rabbit pAb

Cat#: orb771264 (Manual)

For research use only. Not intended for diagnostic use.

Product Name CD88 rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from the N-

terminal region of human C5AR1. AA range:1-50

Specificity CD88 Polyclonal Antibody detects endogenous levels of CD88 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name C5a anaphylatoxin chemotactic receptor

Gene Name C5AR1

Cellular localization Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle.

Phosphorylated C5aR colocalizes with ARRB1 and ARRB2 in cytoplasmic

vesicles. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.





**Clonality** Polyclonal

Concentration 1 mg/ml

**Observed band** 39kD

58503 **Human Gene ID** 

**Human Swiss-Prot Number** P21730


**Alternative Names** C5AR1; C5AR; C5R1; C5a anaphylatoxin chemotactic receptor; C5a-R;

C5aR; ĆD88

**Background** 

function:Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. This receptor stimulates chemotaxis, granule enzyme release and superoxide anion production.,PTM:Sulfation plays a critical role in the association of the receptor with C5a, but no significant role in the ability of the receptor to transduce a signal and mobilize calcium in response to a small peptide agonist., similarity: Belongs to the G-protein coupled

receptor 1 family.,




Western blot analysis of MOUSE-BRAIN RAT-SPLEEN lysis using C5AR1 antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





Explore. Bioreagents.



Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000